


Housatonic Valley Health District

COVID-19 RECOMMENDATIONS: BUSINESSES & MUNICIPALITIES


December 20, 2022



Prepared by: Ruth Quattro, MPI



OUTLINE



Housatonic Valley Health District COVID-19 Recommendations - p. 3 - 4 COVID-19 Vaccination Update - p. 5 - 6 Vaccination Eligibility - p. 7 Definitions - p.8 Quarantine and Self-Isolation Guidelines - p. 9 - 10

HOUSATONIC VALLEY HEALTH DISTRICT



COVID-19 Recommendations

HVHD acknowledges the effectiveness of mask wearing as a way to mitigate the spread of COVID-19.

We respect individuals and businesses choices to unmask.



Wear a well-fitting mask when in crowded groups.



Spend time socializing outside with people who you don't live with, when possible.



Good ventilation can reduce the spread of COVID-19 and lower the risk of exposure.

Enhanced cleaning and **disinfection** is also recommended.

Plan activities to take place **outside** as much as possible.



Get tested if you are
experiencing COVID-19
symptoms or have been notified
of higher-risk close contact
exposure (e.g. household
exposures, crowded social
gatherings).



Practice **proper hygiene**(wash hands regularly,
cover your mouth/nose
when coughing and
sneezing) and stay
home if you are not feel
well.



Self-isolate for <u>5 days</u> if you test positive for COVID-19 & wear a mask around others for an additional 5 days.

Self-isolate for <u>10 days</u> if you are under the age of 3 or unable to wear a mask.

HOUSATONIC VALLEY HEALTH DISTRICT



COVID-19 Recommendations

What Prevention Steps Should You Take Based on Your COVID-19 Community Level?



Stay informed.
Check the CDC's Community
Level and recommendations
frequently.

Low	Medium	High		
 Stay <u>up to date</u> with COVID-19 vaccines <u>Get tested</u> if you have symptoms 	 If you are at high risk for severe illness, talk to your healthcare provider about whether you need to wear a mask and take other precautions Stay up to date with COVID-19 vaccines Get tested if you have symptoms 	 Wear a mask indoors in public Stay up to date with COVID-19 vaccines Get tested if you have symptoms Additional precautions may be needed for people at high risk for severe illness 		
People may choose to mask at any time. People with <u>symptoms, a positive test, or exposure</u> to someone with COVID- 19 should wear a mask. Masks are recommended in indoor public transportation settings and may be required in other places by local or state authorities.				

Source: CDC COVID-19 by County, Retrieved July 18, 2022

COVID-19 Vaccine Eligibility

	Pfizer	Moderna	Johnson & Johnson
Ages 18+	Primary Series: • 2 doses; 3-8 weeks apart Booster: • 1 Bivalent booster; 2 months after the final dose in the primary series	Primary Series: • 2 doses; 4-8 weeks apart Booster: • 1 Bivalent booster; 2 months after the final dose in the primary series	 Primary Series: 1 dose Booster: 1 booster; 2 months after the final dose in the primary series Anyone who received J&J may receive a 2nd dose of Pfizer or Moderna at least 4 months after their 1st booster Ages 50+ can choose to receive a 2nd booster
Ages 12 - 17 years	Primary Series: • 2 doses; 3-8 weeks apart Booster: • 1 Bivalent booster; 2 months after the final dose in the primary series	Primary Series: (ages 6 mo 5 yrs) • 2 doses • 4 - 8 weeks apart Booster: • Not yet authorized	N/A
Ages 5 - 11 years	Primary Series: • 2 doses; 3 - 8 weeks apart Booster: • 1 Monovalent booster; 5 months after the final dose in the primary series	Primary Series: (ages 6 mo 5 yrs) • 2 doses • 4 - 8 weeks apart Booster: • Not yet authorized	N/A
Ages 6 months - 5 years	Primary Series: (ages 6 mo 4 yrs) • 2 doses • 3- 8 weeks apart between 1st & 2nd dose • At least 8 weeks between 2nd & 3rd dose Booster: • 1 Bivalent booster; 8 weeks after second monovalent primary series dose.	Primary Series: (ages 6 mo 5 yrs) • 2 doses • 4 - 8 weeks apart Booster: • 1 Bivalent booster; 2 months after the final dose in the primary series	N/A

Definitions:

- Primary series = 2 3 doses of Pfizer & Moderna; 1 dose of J&J
- Booster = additional dose
- Up to date = you have received all doses in the primary series and booster when eligible.
- Fully vaccinated = 2 weeks after final dose in primary series
- Monovalent = only 1 antigen in a vaccine
- Bivalent = a vaccine that works by stimulating an immune response against 2 different antigens (i.e. omicron BA.4 and BA.5


Schedule an appt:




https://hvhd.us/covid-19/vaccinations/



DEFINITIONS








Keeps someone who is sick or tested positive for COVID-19 without symptoms away from others, even in their own home.

Community Level

COVID-19 Community Levels
help individuals and
communities decide which
prevention actions to take
based on the latest
information and communities
are classified as low, medium,
or high.



Isolation Day Counts:

Day 0 = the day you were tested
Day 1 = the first full day following the day
you were tested

*If you develop symptoms within 10 days of when you were tested, the clock restarts at day 0 on the day of symptom onset

Exposure Day Counts:

Day 0 = the day of your last exposure to someone with COVID-19

Day 1 = the first full day after your last exposure


COVID-19 EXPOSURE RECOMMENDATIONS

Source: CDC: What to Do If You Were Exposed to COVID-19 (updated August 11, 2022)



Key:

Day 0 = the day of your last exposure Day 1 = first full day after your last exposure



ISOLATION: WHEN TO ISOLATE & WHEN TO END ISOLATION

Source: CDC Isolation and Precautions for People with COVID-19 (updated August 11, 2022)

